

University of Groningen

Isolated limb infusion

Kroon, Hidde Maarten

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kroon, H. M. (2009). *Isolated limb infusion*. [Thesis fully internal (DIV), University of Groningen]. [s.n.].

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).


The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Presentations



Presentations

Kroon HM, Lardenoye JHP, Breslau PJ. De incidentie van ongeplande re-operaties in een algemeen heekkundige kliniek als indicator van de kwaliteit van zorg. *Chirurgendagen Nederlandse Vereniging voor Heelkunde, mei 2005, Veldhoven (abstract: Ned Tijdschr Heelkd 2005;14:99-100).*

Kroon HM, Li Q, Whitfield JR, Chang AE. Involvement of Bcl gene family members in anti-4-1BB-induced T-cell survival and anti-tumor reactivity. *Annual Cancer Research Symposium University of Michigan, November 2005, Ann Arbor, USA.*

Kroon HM, Li Q, Whitfield JR, Chang AE. Involvement of Bcl gene family members in anti-4-1BB-induced T-cell survival and anti-tumor reactivity. *Society of Surgical Oncology, March 2006, San Diego, USA (abstract: Ann Surg Oncol 2006;13(S1):67).*

Kroon HM, Lowe L, Cimmino VM, Chang AE, Wong SL, Johnson TM, Sabel MS. Is the 10% rule appropriate for sentinel lymph node biopsy in melanoma? *Society of Surgical Oncology, March 2006, San Diego, USA (abstract: Ann Surg Oncol 2006;13(S1):36).*

Kroon HM, Li Q, Whitfield JR, Chang AE. Anti-41BB-induced T-cell survival and anti-tumor reactivity. *Symposium Experimenteel Onderzoek Heelkundige Specialismen, november 2006, Groningen.*

Moncrieff M, **Kroon HM**, Stalley P, Kam PC, Thompson JF. Outcomes of the use of isolated limb infusion in soft tissue sarcoma of the extremity. *British Association of Plastic, Reconstructive and Aesthetic Surgeons; winter meeting, December 2007, London, England (abstract: BAPRAS abstract book 2007).*

Kroon HM, Moncrieff M, Kam PC, Thompson JF. Regional toxicity after isolated limb infusion with melphalan and actinomycin-D in melanoma patients. *International Congress on Anti-Cancer Treatment, February 2008, Paris, France (abstract: ICACT abstract book 2008;296).*

Kroon HM, Moncrieff M, Kam PC, Thompson JF. Outcomes of isolated limb infusion. A 14-year experience. *Society of Surgical Oncology, March 2008, Chicago, USA (abstract: Ann Surg Oncol 2008;15(S2):10).*

Moncrieff M, **Kroon HM**, Stalley P, Kam PC, Thompson JF. Outcomes of the use of isolated limb infusion in soft tissue sarcoma of the extremity. *Society of Surgical Oncology, March 2008, Chicago, USA (abstract: Ann Surg Oncol 2008;15(S2):27)*.

Kroon HM, Moncrieff M, Kam PCA, Thompson JF. Results of isolated limb infusion for metastatic melanoma. *Royal Australasian College of Surgeons and The College of Surgeons of Hong Kong Conjoint Annual Scientific Congress, May 2008, Hong Kong, China (abstract: ANZ J Surg 2008;78(suppl 1):A50-1)*.

Kroon HM, Moncrieff M, Kam PCA, Thompson JF. The use of isolated limb infusion in advanced soft tissue sarcoma. *Royal Australasian College of Surgeons and The College of Surgeons of Hong Kong Conjoint Annual Scientific Congress, May 2008, Hong Kong, China (abstract: ANZ J Surg 2008;78(suppl 1):A132)*.

Kroon HM, Moncrieff M, Kam PCA, Thompson JF. Regional toxicity after isolated limb infusion with melphalan and actinomycin-D in melanoma patients. *Sydney Cancer Conference, July 2008, Sydney, Australia (abstract: SCC abstract book 2008;148)*.

Kroon HM, Lin DY, Kam PCA, Thompson JF. Safety and efficacy of isolated limb infusion in elderly melanoma patients. *Sydney Cancer Conference, July 2008, Sydney, Australia (abstract: SCC abstract book 2008;149)*.

Kroon HM, Lin DY, Kam PCA, Thompson JF. Efficacy of repeat isolated limb infusion with melphalan and actinomycin-D for recurrent melanoma. *Sydney Cancer Conference, July 2008, Sydney, Australia (abstract: SCC abstract book 2008;150)*.

Kroon HM, Moncrieff M, Kam PCA, Thompson JF. Isolated limb infusion for metastatic melanoma. *Sydney Cancer Conference, July 2008, Sydney, Australia (abstract: SCC abstract book 2008;151)*.

Kroon HM, Kam PC, Thompson JF. Isolated limb infusion for melanoma and sarcoma. *Australasian College of Dermatologists, August 2008, Sydney, Australia*.

Kroon HM, Moncrieff MD, Kam PC, Thompson JF. Isolated limb infusion, a safe and effective alternative to isolated limb perfusion. *Patron's Prize, Royal Prince Alfred Hospital, July 2008, Sydney, Australia*.

Kroon HM, Lin DY, Kam PCA, Thompson JF. Safety and efficacy of isolated limb infusion in elderly patients with advanced locoregional melanoma. *European Society of Surgical Oncology, September 2008, The Hague, the Netherlands (abstract: Eur J Surg Oncol 2008;34:1041).*

Kroon HM, Moncrieff M, Kam PCA, Thompson JF. Determinants of acute regional limb toxicity following isolated limb infusion for melanoma. *European Society of Surgical Oncology, September 2008, The Hague, the Netherlands (abstract: Eur J Surg Oncol 2008;34:1105-6).*

Kroon HM, Lin DY, Kam PCA, Thompson JF. Efficacy of repeat isolated limb infusion with melphalan and actinomycin-D for recurrent melanoma. *European Society of Surgical Oncology, September 2008, The Hague, the Netherlands (abstract: Eur J Surg Oncol 2008;34:1107).*

Kroon HM, Moncrieff M, Kam PC, Thompson JF. Results and limb toxicity after isolated limb infusion for melanoma. *Perspectives in Melanoma XII, October 2008, The Hague, the Netherlands (abstract: Perspectives in Melanoma XII abstract book 2008;287).*

Kroon HM, Lin DY, Kam PC, Thompson JF. Isolated limb infusion in elderly patients with advanced locoregional melanoma. *Perspectives in Melanoma XII, October 2008, The Hague, the Netherlands (abstract: Perspectives in Melanoma XII abstract book 2008;288).*

Kroon HM, Lin DY, Kam PC, Thompson JF. Results of repeat isolated limb infusion for recurrent melanoma. *Clinical Oncological Society of Australia, November 2008, Sydney, Australia (abstract: Asia-Pac J Clin Oncol 2008;4(Suppl 2):A113).*

Kroon HM, Moncrieff M, Kam PC, Thompson JF. Regional limb toxicity following isolated limb infusion for melanoma. *Clinical Oncological Society of Australia, November 2008, Sydney, Australia (abstract: Asia-Pac J Clin Oncol 2008;4(Suppl 2):A114).*

Kroon HM, Lin DY, Kam PC, Thompson JF. Isolated limb infusion in elderly patients with advanced locoregional melanoma. *Clinical Oncological Society of Australia, November 2008, Sydney, Australia (abstract: Asia-Pac J Clin Oncol 2008;4(Suppl 2):A114).*

Kroon HM, Lin DY, Kam PC, Thompson JF. Isolated limb infusion as palliative treatment for advanced limb disease in patients with stage IV melanoma. *Society of Surgical Oncology, March 2009, Phoenix, USA (abstract: Ann Surg Oncol 2009;16(S1):95).*

Kroon HM, Lin DY, Kam PC, Thompson JF. Major amputation for irresectable extremity melanoma after failure of isolated limb infusion. *Society of Surgical Oncology, March 2009, Phoenix, USA (abstract: Ann Surg Oncol 2009;16(S1):100).*

Kroon HM, Lin DY, Kam PC, Thompson JF. Regionale isolatie-infusie bij oudere patiënten met uitgebreide melanoommetastasering van de extremiteiten; veilig en effectief. *Chirurgendagen Nederlandse Vereniging voor Heelkunde, mei 2009, Veldhoven.*

Kroon HM, Lin DY, Kam PC, Thompson JF. Regionale isolatie-infusie bij patiënten met gemetastaseerd melanoom van de extremiteiten: resultaten en toxiciteit. *Chirurgendagen Nederlandse Vereniging voor Heelkunde, mei 2009, Veldhoven.*